

PEDIATRIC IMMUNOLOGY ENROLLMENT FORM

1 PATIENT INFORMATION:

Name: _____
 Address: _____
 City: _____ State: _____ Zip: _____
 Phone: _____ Alt. Phone: _____
 Email: _____
 DOB: _____ Gender: M F Caregiver: _____
 Height: _____ Weight: _____ Allergies: _____

2 PRESCRIBER INFORMATION:

Name: _____
 Address: _____
 City: _____ State: _____ Zip: _____
 Phone: _____ Fax: _____
 NPI: _____ DEA: _____
 Tax I.D.: _____
 Office Contact: _____ Phone: _____

3 STATEMENT OF MEDICAL NECESSITY: (Please Attach All Medical Documentation)

Date of Diagnosis: _____ Patient also taking Methotrexate? Yes No
 ICD-10: _____ Other: _____ Serious or active infection present? Yes No
 TB Test: Positive Negative Date: _____ Hep B and Hep C ruled out or
 LFT: ALT: _____ AST: _____ Date: _____ treatment started? Yes No
 Assessment: Moderate Mod to Severe Severe Does patient have latex allergy? Yes No
 _____% BSA affected
 Scalp Face Chest Arms Hands Nails
 Back Groin Buttocks Legs Other: _____
 ISGA or EASI

If Prior Authorization is denied, recommended formulary alternatives will be provided to the prescriber based upon the patient's insurance coverage.

Prior Failed Treatments:	Indicate Drug Name and Length of Treatment:
<input type="checkbox"/> 5-ASA	_____
<input type="checkbox"/> Biologics	_____
<input type="checkbox"/> Corticosteroids	_____
<input type="checkbox"/> Immunosuppressants	_____
<input type="checkbox"/> Methotrexate	_____
<input type="checkbox"/> NSAIDS	_____
<input type="checkbox"/> Surgery	_____
<input type="checkbox"/> Topical/Oral Antibiotics	_____
<input type="checkbox"/> UVA <input type="checkbox"/> UVB	_____
<input type="checkbox"/> Others	_____

4 INJECTION TRAINING: To Be Administered by Pharmacist (State of Missouri Only) Pharmacist to Provide Training Patient Trained in MD Office Manufacturer Nurse Support

5 PICK UP OR DELIVERY: Delivery to Patient's Home Delivery to Physician's Office Pharmacy to Coordinate

6 INSURANCE INFORMATION: Please Include Front and Back Copies of Pharmacy and Medical Card

PRESCRIPTION INFORMATION: (Please be sure to choose both induction and maintenance dose where applicable)

Patient Name: _____ Patient's Date of Birth: _____

Medication	Dosage & Strength	Direction	QTY	Refills
<input type="checkbox"/> AMJEVITA™	<input type="checkbox"/> 10mg/0.2ml Prefilled Syringe <input type="checkbox"/> 20mg/0.4ml Prefilled Syringe <input type="checkbox"/> 40mg/0.8ml Prefilled Syringe <input type="checkbox"/> 40mg/0.8ml Pen	Juvenile Idiopathic Arthritis (2 years of age and older) <input type="checkbox"/> 10kg to <15kg: Inject 10mg every other week <input type="checkbox"/> 15kg to <30kg: Inject 20mg every other week <input type="checkbox"/> ≥30kg: Inject 40mg every other week	2	
		Crohn's Disease (6 years of age and older): Induction Dose: <input type="checkbox"/> 17kg to <40kg: Inject 80mg on day 1 and 40mg day 15 <input type="checkbox"/> ≥40kg: Inject 160mg (single-dose or split over two consecutive days) on day 1 and 80mg day 15	3	
		Maintenance Dose: <input type="checkbox"/> 17kg to <40kg: Inject 80mg on day 1 and 40mg day 15 <input type="checkbox"/> ≥40kg: Inject 160mg (single-dose or split over two consecutive days) on day 1 and 80mg day 15	6	
		PsO pediatric patients 6 years and older: <input type="checkbox"/> <50kg: 75mg at week 0, 1, 2, 3, and 4 and every 4 weeks thereafter <input type="checkbox"/> ≥50kg: 150mg at week 0, 1, 2, 3, and 4 and every 4 weeks thereafter	3	
<input type="checkbox"/> COSENTYX®	<input type="checkbox"/> 150mg/ml single-dose Sensoready® Pen <input type="checkbox"/> 150mg/ml single-dose Prefilled Syringe <input type="checkbox"/> 75mg/0.5ml single-dose Prefilled Syringe <input type="checkbox"/> 300mg/2ml single-dose UnoReady pen <input type="checkbox"/> 300mg/2ml single-dose prefilled Syringe <input type="checkbox"/> For Healthcare Professional Use Only: 150mg lyophilized powder in a single-dose vial for reconstitution	PsA pediatric patients 2 years and older: <input type="checkbox"/> ≥15kg and <50kg: 75mg at week 0, 1, 2, 3, and 4 and every 4 weeks thereafter <input type="checkbox"/> ≥50kg: 150mg at week 0, 1, 2, 3, and 4 and every 4 weeks thereafter	6	
		Enthesitis-Related Arthritis 4 years and older: <input type="checkbox"/> ≥15kg and <50kg: 75mg at weeks 0, 1, 2, 3, and 4 and every 4 weeks thereafter <input type="checkbox"/> ≥50kg: 150mg at week 0, 1, 2, 3, and 4 and every 4 weeks thereafter	6	
<input type="checkbox"/> DUPIXENT®	Pediatric Atopic Dermatitis (6 months to 17 years of age): <input type="checkbox"/> 300mg/2ml single-dose Prefilled Syringe <input type="checkbox"/> 200mg/1.14ml single-dose Prefilled Syringe <input type="checkbox"/> 300mg/2ml single-dose Prefilled Pen <input type="checkbox"/> 200mg/1.14ml single-dose Prefilled Pen	Induction Dose: <input type="checkbox"/> ≥60kg: Inject 600mg SC (two 300mg injections) <input type="checkbox"/> 30 to <60kg: Inject 400mg SC (two 200mg injections) <input type="checkbox"/> 15 to <30kg: Inject 600mg SC (two 300mg injections)	2	0
		Maintenance Dose: <input type="checkbox"/> ≥60kg: Inject 300mg SC every other week <input type="checkbox"/> 30 to <60kg: Inject 200mg SC every other week <input type="checkbox"/> 15 to <30kg: Inject 300mg SC every 4 weeks		
<input type="checkbox"/> EUCRISA®	Pediatric patients 3 months of age and older: <input type="checkbox"/> 2% Ointment	Induction Dose: <input type="checkbox"/> ≥60kg: Inject 600mg SC (two 300mg injections) <input type="checkbox"/> 30 to <60kg: Inject 400mg SC (two 200mg injections) <input type="checkbox"/> 15 to <30kg: Inject 600mg SC (two 300mg injections)	60g	0
		Maintenance Dose: <input type="checkbox"/> ≥60kg: Inject 300mg SC every other week <input type="checkbox"/> 30 to <60kg: Inject 200mg SC every other week <input type="checkbox"/> 15 to <30kg: Inject 300mg SC every 4 weeks	100g	
<input type="checkbox"/> ENBREL®	<input type="checkbox"/> 50mg/ml Sureclick® Autoinjector <input type="checkbox"/> 50mg/ml Enbrel Min® Prefilled Cartridge For Enbrel Mini® only: AutoTouch® reusable Autoinjector <input type="checkbox"/> 50mg/ml Prefilled Syringe <input type="checkbox"/> 25mg/0.5ml Prefilled Syringe <input type="checkbox"/> 25mg Lyophilized Power Vial <input type="checkbox"/> Other: _____	Induction Dose: <input type="checkbox"/> ≥63kg: Inject 50mg weekly <input type="checkbox"/> <63kg: Inject 0.8mg/kg weekly (maximum 50mg/week) (To achieve pediatric doses other than 50mg or 25mg, use reconstituted Enbrel lyophilized powder)		
		Maintenance Dose: <input type="checkbox"/> ≥63kg: Inject 50mg weekly <input type="checkbox"/> <63kg: Inject 0.8mg/kg weekly (maximum 50mg/week) (To achieve pediatric doses other than 50mg or 25mg, use reconstituted Enbrel lyophilized powder)		

PRESCRIBER SIGNATURE: I authorize pharmacy to act as my designee for initiating and coordinating insurance prior authorizations, nursing services and patient assistance programs.

Signature: _____ Date: _____ Signature: _____ Date: _____

Substitution Permitted

Dispense As Written

Prior authorization approval and insurance benefits will be determined by the payor based upon the patient's eligibility, medical necessity, and the terms of the patient's coverage, among other things. Participation in this program is not a guarantee of prior authorization or of payment.

Confidentiality Notice: This fax is intended to be delivered only to the named addressee and contains confidential information that may be protected health information under federal and state laws. If you are not the named addressee, you should not disseminate, distribute or copy this fax. Please inform the sender immediately if you have received this document in error and then destroy this document immediately.

PEDIATRIC IMMUNOLOGY ENROLLMENT FORM

Lorem ipsum



1 PATIENT INFORMATION:

Name: _____
 Address: _____
 City: _____ State: _____ Zip: _____
 Phone: _____ Alt. Phone: _____
 Email: _____
 DOB: _____ Gender: M F Caregiver: _____
 Height: _____ Weight: _____ Allergies: _____

2 PRESCRIBER INFORMATION:

Name: _____
 Address: _____
 City: _____ State: _____ Zip: _____
 Phone: _____ Fax: _____
 NPI: _____ DEA: _____
 Tax I.D.: _____
 Office Contact: _____ Phone: _____

3 STATEMENT OF MEDICAL NECESSITY: (Please Attach All Medical Documentation)

Date of Diagnosis: _____ Patient also taking Methotrexate? Yes No
 ICD-10: _____ Other: _____ Serious or active infection present? Yes No
 TB Test: Positive Negative Date: _____ Hep B and Hep C ruled out or
 LFT: ALT: _____ AST: _____ Date: _____ treatment started? Yes No
 Assessment: Moderate Mod to Severe Severe Does patient have latex allergy? Yes No
 _____% BSA affected
 Scalp Face Chest Arms Hands Nails
 Back Groin Buttocks Legs Other: _____
 ISGA or EASI

If Prior Authorization is denied, recommended formulary alternatives will be provided to the prescriber based upon the patient's insurance coverage.

Prior Failed Treatments:	Indicate Drug Name and Length of Treatment:
<input type="checkbox"/> 5-ASA	_____
<input type="checkbox"/> Biologics	_____
<input type="checkbox"/> Corticosteroids	_____
<input type="checkbox"/> Immunosuppressants	_____
<input type="checkbox"/> Methotrexate	_____
<input type="checkbox"/> NSAIDS	_____
<input type="checkbox"/> Surgery	_____
<input type="checkbox"/> Topical/Oral Antibiotics	_____
<input type="checkbox"/> UVA <input type="checkbox"/> UVB	_____
<input type="checkbox"/> Others	_____

4 INJECTION TRAINING: To Be Administered by Pharmacist (State of Missouri Only) Pharmacist to Provide Training Patient Trained in MD Office Manufacturer Nurse Support

5 PICK UP OR DELIVERY: Delivery to Patient's Home Delivery to Physician's Office Pharmacy to Coordinate

6 INSURANCE INFORMATION: Please Include Front and Back Copies of Pharmacy and Medical Card

PRESCRIPTION INFORMATION: (Please be sure to choose both induction and maintenance dose where applicable)

Patient Name: _____ Patient's Date of Birth: _____

Medication	Dosage & Strength	Direction	QTY	Refills
<input type="checkbox"/> HUMIRA®	Hidradenitis Suppurativa <input type="checkbox"/> Adolescent Hidradenitis Suppurativa 80mg/0.8ml and 40mg/0.4ml Starter pack <input type="checkbox"/> Adolescent Hidradenitis Suppurativa 40mg/0.4ml Starter Package <input type="checkbox"/> Adolescent Hidradenitis Suppurativa 40mg/0.8ml Starter Package <input type="checkbox"/> 40mg/0.4ml Pen <input type="checkbox"/> 40mg/0.8ml Pen <input type="checkbox"/> 80mg/0.8ml Pen <input type="checkbox"/> 40mg/0.4ml Prefilled Syringe <input type="checkbox"/> 40mg/0.8ml Prefilled Syringe <input type="checkbox"/> 80mg/0.8ml Prefilled Syringe	<input type="checkbox"/> Induction Dose (Adolescents 12 years and older): <input type="checkbox"/> 30kg to <60kg: Inject 80mg SC on day 1, then 40mg SC on day 8 and every other week thereafter <input type="checkbox"/> ≥60kg: Inject two 80mg pens SC on day 1, then one 80mg pen SC on day 15 <input type="checkbox"/> ≥60kg: Inject one 80mg pen SC on day 1, then 80mg pen SC on day 2, then one 80mg pen SC on day 15 <input type="checkbox"/> Maintenance Dose (Adolescents 12 years and older): <input type="checkbox"/> 30kg to <60kg: Inject 40mg every other week <input type="checkbox"/> ≥60kg: Inject 40mg on day 29 then Inject 40mg every week <input type="checkbox"/> ≥60kg: Inject 80mg on day 29 then Inject 80mg every other week		
<input type="checkbox"/> HUMIRA®	Juvenile Idiopathic Arthritis + Pediatric Uveitis <input type="checkbox"/> 10mg/0.1ml Prefilled Syringe <input type="checkbox"/> 10mg/0.2ml Prefilled Syringe <input type="checkbox"/> 20mg/0.2ml Prefilled Syringe <input type="checkbox"/> 20mg/0.4ml Prefilled Syringe <input type="checkbox"/> 40mg/0.4ml Prefilled Syringe <input type="checkbox"/> 40mg/0.8ml Prefilled Syringe <input type="checkbox"/> 40mg/0.4ml Pen <input type="checkbox"/> 40mg/0.8ml Pen	<input type="checkbox"/> Weight-Based Dosing (5 years and older): <input type="checkbox"/> 10kg to <15kg: Inject 10mg SC every other week <input type="checkbox"/> 15kg to <30kg: Inject 20mg SC every other week <input type="checkbox"/> ≥30kg: Inject 40mg SC every other week		
<input type="checkbox"/> HUMIRA®	Pediatric Crohn's Disease <input type="checkbox"/> Pediatric Crohn's Starter Pack Prefilled Syringe: <input type="checkbox"/> 80mg/0.8ml, 40mg/0.4ml <input type="checkbox"/> 40mg/0.8ml <input type="checkbox"/> 80mg/0.8ml <input type="checkbox"/> Maintenance pack <input type="checkbox"/> 20mg/0.2ml Prefilled Syringe <input type="checkbox"/> 20mg/0.4ml Prefilled Syringe <input type="checkbox"/> 40mg/0.4ml Pen <input type="checkbox"/> 40mg/0.4ml Prefilled Syringe	<input type="checkbox"/> Induction Dose (Pediatrics patients 6 years and older): <input type="checkbox"/> 17kg to <40kg: Inject one 80mg pen SC on day 1, then one 40mg pen SC on day 15 <input type="checkbox"/> ≥40kg: Inject two 80mg pens SC on day 1, then one 80mg pen SC on day 15 <input type="checkbox"/> ≥40kg: Inject one 80mg pen SC on day 1, then 80mg pen SC on day 2, then one 80mg pen SC on day 15 <input type="checkbox"/> Maintenance Dose (Pediatrics patients 6 years and older): <input type="checkbox"/> 17kg to <40kg: Inject 20mg SC every other week <input type="checkbox"/> ≥40kg: Inject 40mg SC every other week	2 3	0 0
<input type="checkbox"/> HUMIRA®	Pediatric Ulcerative Colitis <input type="checkbox"/> Pediatric Ulcerative Colitis' Starter Pack: 80mg/0.8ml Pen <input type="checkbox"/> 20mg/0.2ml Prefilled Syringe <input type="checkbox"/> 20mg/0.4ml Prefilled Syringe <input type="checkbox"/> 40mg/0.4ml Pen <input type="checkbox"/> 40mg/0.8ml Pen <input type="checkbox"/> 80mg/0.8ml Pen <input type="checkbox"/> 40mg/0.4ml Prefilled Syringe <input type="checkbox"/> 40mg/0.8ml Prefilled Syringe <input type="checkbox"/> 80mg/0.8ml Prefilled Syringe	<input type="checkbox"/> Induction Dose (5 years and older): <input type="checkbox"/> 20kg to 40kg: Inject 80mg SC at week 0 (day 1), then 40mg SC at week 1 (day 8), then 40mg SC at week 2 (day 15) <input type="checkbox"/> >40kg: Inject 160mg SC at week 0 (day 1), then 80mg SC at week 1 (day 8), then 40mg SC at week 2 (day 15) <input type="checkbox"/> Maintenance Dose (Pediatric patients 5 years and older): <input type="checkbox"/> 20kg to 40kg: Inject 40mg SC at week 4 (day 29) and every other week thereafter <input type="checkbox"/> 20kg to 40kg: Inject 20mg SC at week 4 (day 29) and every other week thereafter <input type="checkbox"/> 40kg: Inject 80mg SC at week 4 (day 29) and every other week thereafter <input type="checkbox"/> 40kg: Inject 40mg SC at week 4 (day 29) and every other week thereafter		
<input type="checkbox"/>				

All strengths and dosages listed are Humira® Citrate Free

PRESCRIBER SIGNATURE: I authorize pharmacy to act as my designee for initiating and coordinating insurance prior authorizations, nursing services and patient assistance programs.

Signature: _____ Date: _____ Signature: _____ Date: _____

Substitution Permitted

Dispense As Written

Prior authorization approval and insurance benefits will be determined by the payor based upon the patient's eligibility, medical necessity, and the terms of the patient's coverage, among other things. Participation in this program is not a guarantee of prior authorization or of payment.

Confidentiality Notice: This fax is intended to be delivered only to the named addressee and contains confidential information that may be protected health information under federal and state laws. If you are not the named addressee, you should not disseminate, distribute or copy this fax. Please inform the sender immediately if you have received this document in error and then destroy this document immediately.

v10.3_062623

©2020 KloudScript, Inc. - All rights reserved.

PEDIATRIC IMMUNOLOGY ENROLLMENT FORM

1 PATIENT INFORMATION:

Name: _____
 Address: _____
 City: _____ State: _____ Zip: _____
 Phone: _____ Alt. Phone: _____
 Email: _____
 DOB: _____ Gender: M F Caregiver: _____
 Height: _____ Weight: _____ Allergies: _____

2 PRESCRIBER INFORMATION:

Name: _____
 Address: _____
 City: _____ State: _____ Zip: _____
 Phone: _____ Fax: _____
 NPI: _____ DEA: _____
 Tax I.D.: _____
 Office Contact: _____ Phone: _____

3 STATEMENT OF MEDICAL NECESSITY: (Please Attach All Medical Documentation)

Date of Diagnosis: _____ Patient also taking Methotrexate? Yes No
 ICD-10: _____ Other: _____ Serious or active infection present? Yes No
 TB Test: Positive Negative Date: _____ Hep B and Hep C ruled out or
 LFT: ALT: _____ AST: _____ Date: _____ treatment started? Yes No
 Assessment: Moderate Mod to Severe Severe Does patient have latex allergy? Yes No
 _____% BSA affected
 Scalp Face Chest Arms Hands Nails
 Back Groin Buttocks Legs Other: _____
 ISGA or EASI _____

If Prior Authorization is denied, recommended formulary alternatives will be provided to the prescriber based upon the patient's insurance coverage.

Prior Failed Treatments:	Indicate Drug Name and Length of Treatment:
<input type="checkbox"/> 5-ASA	_____
<input type="checkbox"/> Biologics	_____
<input type="checkbox"/> Corticosteroids	_____
<input type="checkbox"/> Immunosuppressants	_____
<input type="checkbox"/> Methotrexate	_____
<input type="checkbox"/> NSAIDS	_____
<input type="checkbox"/> Surgery	_____
<input type="checkbox"/> Topical/Oral Antibiotics	_____
<input type="checkbox"/> UVA <input type="checkbox"/> UVB	_____
<input type="checkbox"/> Others	_____

4 INJECTION TRAINING: To Be Administered by Pharmacist (State of Missouri Only) Pharmacist to Provide Training Patient Trained in MD Office Manufacturer Nurse Support

5 PICK UP OR DELIVERY: Delivery to Patient's Home Delivery to Physician's Office Pharmacy to Coordinate

6 INSURANCE INFORMATION: Please Include Front and Back Copies of Pharmacy and Medical Card

PRESCRIPTION INFORMATION: (Please be sure to choose both induction and maintenance dose where applicable)

Patient Name: _____ Patient's Date of Birth: _____

Medication	Dosage & Strength	Direction	QTY	Refills
<input type="checkbox"/> HUMIRA® Biosimilar	<input type="checkbox"/> _____ <input type="checkbox"/> _____			
<input type="checkbox"/> OPZELURA™	12 years of age and older: <input type="checkbox"/> 1.5% Cream	<input type="checkbox"/> Apply a thin layer twice daily to affected areas <i>Do not use more than 60 grams per week</i>	60g	
<input type="checkbox"/> RINVOQ® XR	<input type="checkbox"/> 15mg ER Tablet <input type="checkbox"/> 30mg ER Tablet	<input type="checkbox"/> Pediatric patients 12 years of age and older weighing at least 40kg: <input type="checkbox"/> Take 15mg tablet once daily "If an adequate response is not achieved, consider increasing the dosage to 30mg orally once daily"	30	
<input type="checkbox"/> STELARA®	<input type="checkbox"/> 45mg/0.5ml Prefilled Syringe <input type="checkbox"/> 45mg/0.5ml Single-Dose Vial <input type="checkbox"/> 90mg/ml Prefilled Syringe	Psoriasis Pediatric Dose (6 to 17 years old): <input type="checkbox"/> <60kg: Inject 0.75mg/kg SC initially at 4 weeks, then every 12 weeks thereafter <input type="checkbox"/> 60kg to 100kg: Inject 45mg SC initially at 4 weeks, then every 12 weeks thereafter <input type="checkbox"/> >100kg: Inject 90mg SC initially at 4 weeks, then every 12 weeks thereafter	1	0
		Psoriatic Arthritis Pediatric Dose (6 to 17 years old): <input type="checkbox"/> <60kg: Inject 0.75mg/kg SC initially at 4 weeks, then every 12 weeks thereafter <input type="checkbox"/> >60kg: Inject 45mg SC initially at 4 weeks, then every 12 weeks thereafter <input type="checkbox"/> >100kg with co-existent moderate-to-severe plaque psoriasis: Inject 90mg SC initially at 4 weeks, then every 12 weeks thereafter		0
<input type="checkbox"/> TALTZ®	Pediatric Plaque Psoriasis <input type="checkbox"/> 80mg/ml Single-Dose Prefilled Autoinjector <input type="checkbox"/> 80mg/ml Single-Dose Prefilled Syringe <i>20mg and 40mg doses for patients weighing ≤50kg must be prepared and administered by a qualified healthcare professional.</i>	Induction Dose (6 years and older): <input type="checkbox"/> >50kg: Inject 160mg SC (two 80mg injections) at week 0 <input type="checkbox"/> 25 to 50kg: Inject 80 mg SC at week 0 <input type="checkbox"/> <25kg: Inject 40mg SC at week 0	2	0
		Maintenance Dose (6 years and older): <input type="checkbox"/> >50kg: Inject 80mg SC at week 4 and every 4 weeks thereafter <input type="checkbox"/> 25 to 50kg: Inject 40 mg SC at week 4 and every 4 weeks thereafter <input type="checkbox"/> <25kg: Inject 20 mg at week 4 and every 4 weeks thereafter	1	
<input type="checkbox"/> XELJANZ®	Polyarticular Course Juvenile Idiopathic Arthritis (pcJIA) <input type="checkbox"/> 1mg/ml oral solution <input type="checkbox"/> 5mg tablets	Weight-Based Dosing (Pediatric patients 2 years and older): <input type="checkbox"/> 10kg to <20kg: Take 3.2mg (3.2ml oral solution) twice daily <input type="checkbox"/> 20kg to <40kg: Take 4mg (4ml oral solution) twice daily <input type="checkbox"/> ≥40kg: Take 5mg (one 5mg oral tablet or 5ml oral solution) twice daily <input type="checkbox"/> Take 5mg by mouth twice daily	60	
<input type="checkbox"/> YUSIMRY™	Juvenile Idiopathic Arthritis + Pediatric Crohn's Disease <input type="checkbox"/> 40mg/0.8ml Prefilled Syringe <input type="checkbox"/> 40mg/0.8ml Prefilled Pen	Juvenile idiopathic arthritis pediatric patients 2 years of age and older >30kg: <input type="checkbox"/> Inject 40mg SC every other week	2	
		CD pediatric patients 6 years of age and older >40kg: <input type="checkbox"/> Induction Dose: Inject 160mg SC on day 1 and 80mg on day 15 <input type="checkbox"/> Maintenance Dose: Inject 40mg SC every other week (Starting on Day 29)	6	2
<input type="checkbox"/> _____				

PRESCRIBER SIGNATURE: I authorize pharmacy to act as my designee for initiating and coordinating insurance prior authorizations, nursing services and patient assistance programs.

Signature: _____ Date: _____ Signature: _____ Date: _____

Substitution Permitted

Dispense As Written

Prior authorization approval and insurance benefits will be determined by the payor based upon the patient's eligibility, medical necessity, and the terms of the patient's coverage, among other things. Participation in this program is not a guarantee of prior authorization or of payment.

Confidentiality Notice: This fax is intended to be delivered only to the named addressee and contains confidential information that may be protected health information under federal and state laws. If you are not the named addressee, you should not disseminate, distribute or copy this fax. Please inform the sender immediately if you have received this document in error and then destroy this document immediately.