

## RHEUMATOID ARTHRITIS SPECIALTY CARE PROGRAM

Phone: **833-796-6470** • Fax: **844-841-3401** 



PATIENT INFORMATION:  Name:		PRESCRIBER INFORMATION: Name:		
		A -I -I		
tv:	State: Zip:	City: State:	Zin:	
one:	Alt. Phone:	Phone: Fax:	_ <del>-</del> - P·	
nail:		NPI: DEA:		
B: Ge	ender: O M O F Caregiver:			
ight: We	eight: Allergies:	Tax I.D.: Phone:		
	OF MEDICAL NECESSITY: (Please Att.	ach All Medical Documentation) Prior Failed Trea		
te of Diagnosis:	Patient also taking Method	☐ Azulfidine® ☐ Celebrex® trexate? ☐ Yes ☐ No ☐ Biologics ☐ Corticoste	atments:	hotrex
7_10.	Serious or active infection	present?  Yes No Calcipotriene Indocin®		
7-10	Serious or active infection  Hep B ruled out or treatments	ent started?  Yes No Indicate Drug Name and Le	ength of Treati	ment:
	Does patient have latex at	liergy? Li Yes Li No		
	egative Date: LFT: ALT: AST:_ denied, recommended formulary alternatives	Date:  will be provided to the prescriber based upon the patient's	s insurance co	verag
	<del>-</del>	Missouri Only) O Pharmacist to Provide Training O Patient Trained in MD Office O		
		ome O Delivery to Physician's Office O Pharma		nate
		and Back Copies of Pharmacy and Medical Card		
tient Name:	INFORMATION: (Please be sure to c	choose both induction and maintenance dose wh	ere applica	ble)
Medication	Dosage & Strength	Direction	QTY	Refil
□ ACTEMRA®	☐ 162mg/0.9ml Prefilled Syringe	☐ Inject 162mg SC every other week (< 220 lbs) ☐ Inject 162mg SC every week (> 220 lbs) ☐ Inject 162mg SC every 2 weeks (> 66 lbs) ☐ Inject 162mg SC every 3 weeks (< 66 lbs)		
□ CIMZIA®	☐ Prefilled Syringe Starter Kit☐ 200mg/ml Prefilled Syringe	☐ Induction Dose: Inject 400mg SC on day 1, day 14 and day 28	6	0
u CiiviziA	☐ 200mg Lyophilized Powder Vial	☐ Maintenance: Inject 400mg SC every 4 weeks ☐ Maintenance: Inject 200mg SC every other week	2	
□ COSENTYX®	☐ 150mg/ml Sensoready® Pen	☐ Induction Dose: Inject 150mg SC at weeks 0, 1, 2, 3, and 4	5	0
	☐ 150mg/ml Prefilled Syringe☐ 150mg/ml Lyophilized Powder Vial	☐ Induction Dose: Inject 300mg SC at weeks 0, 1, 2, 3, and 4☐ Maintenance Dose: Inject 150mg SC every four weeks	10	0
	☐ 50mg/ml Sureclick Autoinjector	☐ Maintenance Dose: Inject 300mg SC every four weeks	2	
	□ 50mg/ml Enbrel Mini™ Prefilled Cartridge	☐ Inject 50mg SC once a week	4	
□ ENBREL®	<ul> <li>☐ For Enbrel Mini™ only: AutoTouch™ Autoinjector</li> <li>☐ 50mg/ml Prefilled Syringe</li> </ul>	☐ Inject 25mg SC twice a week (3-4 days apart) ☐ Other	1	
	☐ 25mg/ml Prefilled Syringe☐ 25mg/ml Vial	U Otner		
	☐ 40mg/0.4ml Pen			
□ HUMIRA®	<ul><li>□ 40mg/0.4ml Prefilled Syringe</li><li>□ 40mg/0.8ml Pen</li></ul>	☐ Inject 40mg SC every other week ☐ Patient has signed ☐ HUMIRA Complete form		
	□ 40mg/0.8ml Prefilled Syringe	,		
	All strengths and dosage ☐ 150mg/1.14ml Prefilled Syringe	ges listed are Humira® Citrate Free		
□ KEVZARA®	☐ 150mg/1.14ml Prefilled Pen☐ 200mg/1.14ml Prefilled Syringe☐ 200mg/1.14ml Prefilled Pen☐ 200mg/1.14ml Pen☐ 200mg/1	□ Inject 150mg SC every 2 weeks	2	
	□ 200mg/1.14ml Prefilled Pen	☐ Inject 200mg SC every 2 weeks	2	
□ ORENCIA®	<ul> <li>□ 250mg Lyophilized Powder Vial</li> <li>□ 125mg/ml ClickJect™ Autoinjector</li> <li>□ 50mg/0.4ml Prefilled Syringe</li> <li>□ 87.5mg/0.7ml Prefilled Syringe</li> <li>□ 125mg/ml Prefilled Syringe</li> </ul>	☐ Induction Dose: Patient Weight < 132 lbs: 500mg; 132-220 lbs: 750mg; > 220 lbs: 1000mg administered IV, then inject 125mg SC within 24 hours		0
UNENUA"		☐ Inject 50mg SC once a week (10 to less than 25kg)☐ Inject 87.5mg SC once a week (25 to less than 50kg)	4 4	
		☐ Inject 125mg SC once a week (50kg or more)	4	
OTEZLA® (for PsA)	☐ Starter Pack (Titration) ☐ 30mg Tablets	☐ Starter Pack: Take one tablet in the morning on day 1, then take one tablet in the morning and one tablet in the evening as	1	0
		directed on the starter pack  Maintenance: Take one 30mg tablet by mouth twice daily	60	
□ SIMPONI®	☐ 50mg/0.5ml Smartject Autoinjector☐ 50mg/0.5ml Prefilled Syringe	☐ Inject 50mg SC once a month	1	
	☐ 45mg/0.5ml Prefilled Syringe (for < 220 lbs)	☐ Induction Dose: Inject 1 prefilled syringe SC on day 1	1	
	☐ 90mg/1ml Prefilled Syringe (for > 220 lbs)	☐ Maintenance: Inject 1 prefilled syringe SC on day 29, and every 12 weeks thereafter sthat patient has been trained and is eligible for self-injection	1	0
(for PsA)	Error Toblet	☐ Take one 5mg tablet by mouth twice a day	60	
(for PsA)  XELJANZ®	□ 5mg Tablet	B = 1		
(for PsA)  XELJANZ®  XELJANZ® XR	☐ 11mg Tablet	☐ Take one 11mg tablet once a day	30	
☐ XELJANZ® ☐ XELJANZ® XR ☐ RASUVO®	☐ 11mg Tablet	•		
(for PsA)  XELJANZ®  XELJANZ® XR	☐ 11mg Tablet	,	30	