

HYPERCHOLESTEROLEMIA SPECIALTY CARE PROGRAM

Phone: 833-796-6470 • Fax: 844-841-3401



PATIENT INFORMATION: Name:		PRESCRIBER INFORMATION: Name:			
	State: Zip:	City:	State:	Zip:	
-	Alt. Phone:	Phone:	Fax:		
		NPI:	DEA:		
		14X 1.D			
DOB:	Gender: O M O F Caregiver:	Office Contact	ct: Phone:		
-leight:	Weight: Allergies:	Specialty: □	l Cardiology □ Lipidology □	Other	
3 STATEMEN	T OF MEDICAL NECESSITY: (Please	e Attach All Medic	al Documentation and Laborato	ry Results)	
Date of Diagnosis: _			Prior Indica	te Drug Name	
Primary ICD-10:	Secondary ICD-10:		Failed Therapies: and Leng	th of Treatment:	
Other:			☐ Fibrates		
Contraindications:			☐ Niacin ☐ Omega-3		
Fibrates: 🗆 Yes 🗅 No Statin: 🗅 Yes 🗅 No Niacin: 🗅 Yes 🗅 No			☐ Statin		
If yes: ☐ Myopathy or Rhabdomyolysis ☐ Hepatic Disease ☐ Renal Dysfunc☐ Pregnancy or Lactation ☐ Recent Stroke or TIA ☐ Other					
	ctation - Recent Stroke of HA - Other		Other		
Laboratory Tests: ☐ Lipid Panel	□ No □ Yes Date:		If Prior Authorization is denied,		
Liver Function			formulary alternatives will be provided to the prescriber based upon the patient's insurance		
■ Renal Function			coverage.		
f labs must be obta	ined from another prescriber, please indicate na	ame here:			
1 INJECTION	TRAINING: O To Be Administered by Pharmacist	Pharmacist to Provide Tra	aining O Patient Trained in MD Office O M	anufacturer Nurse Support	
	R DELIVERY: O Delivery to Patient's Ho				
	E INFORMATION: Please Include Front				
	N INFORMATION:				
			Patient's Date of Birth:		
Medication	Dosage & Strength		Direction	QTY Refills	
	☐ 75mg/ml single-dose Prefilled Autoinjector/Pen☐ 150mg/ml single-dose Prefilled Autoinjector/Pen	☐ 75mg SQ once every 2 weeks ☐ 300mg (150mg/2pen) SQ once every 4 weeks In adults with HeFH undergoing LDL apheresis or in adults with HoFH:			
☐ PRALUENT®				2 l	
	,	☐ 150mg SQ once ev			
		☐ 140mg SQ every 2		emia:	
	140ma/ml single deep Punfilled Curriers	☐ 140mg SQ every 2 ☐ 420mg SQ once m	weeks onthly	emia:	
□ pedatua®	☐ 140mg/ml single-dose Prefilled Syringe☐ 140mg/ml single-dose Prefilled SureClick®	☐ 140mg SQ every 2 ☐ 420mg SQ once m In pediatric patients a ☐ 140mg SQ every 2	weeks onthly aged 10 years and older with HeFH: weeks	emia:	
☐ REPATHA®	☐ 140mg/ml single-dose Prefilled SureClick® Autoinjector	☐ 140mg SQ every 2☐ 420mg SQ once m In pediatric patients a☐ 140mg SQ every 2☐ 420mg SQ once m	weeks onthly aged 10 years and older with HeFH: weeks onthly		
☐ REPATHA®	☐ 140mg/ml single-dose Prefilled SureClick®	☐ 140mg SQ every 2☐ 420mg SQ once m In pediatric patients a☐ 140mg SQ every 2☐ 420mg SQ once m	weeks onthly aged 10 years and older with HeFH: weeks onthly ic patients aged 10 years and older with Ho		
☐ REPATHA®	☐ 140mg/ml single-dose Prefilled SureClick® Autoinjector	□ 140mg SQ every 2 □ 420mg SQ once m In pediatric patients: □ 140mg SQ every 2 □ 420mg SQ once m In adults and pediatri □ 420mg SQ once m □ 420mg SQ once m	weeks onthly aged 10 years and older with HeFH: weeks onthly c patients aged 10 years and older with Ho onthly weeks		
☐ REPATHA®	☐ 140mg/ml single-dose Prefilled SureClick® Autoinjector	□ 140mg SQ every 2 □ 420mg SQ once m In pediatric patients: □ 140mg SQ every 2 □ 420mg SQ once m In adults and pediatri □ 420mg SQ once m □ 420mg SQ once m	weeks onthly aged 10 years and older with HeFH: weeks onthly ic patients aged 10 years and older with Ho onthly		
☐ REPATHA® ☐ OTHER	☐ 140mg/ml single-dose Prefilled SureClick® Autoinjector	I 140mg SQ every 2 420mg SQ once m In pediatric patients: 140mg SQ every 2 420mg SQ once m In adults and pediatri 420mg SQ once m 420mg SQ once m 420mg SQ every 2 (For 420mg: Inject the	weeks onthly aged 10 years and older with HeFH: weeks onthly c patients aged 10 years and older with Ho onthly weeks		
OTHER	☐ 140mg/ml single-dose Prefilled SureClick® Autoinjector ☐ 420mg/3.5ml single-dose Pushtronex® system	□ 140mg SQ every 2 □ 420mg SQ once m In pediatric patients: □ 140mg SQ every 2 □ 420mg SQ once m In adults and pediatri □ 420mg SQ once m □ 420mg SQ oevery 2 (For 420mg: Inject the within 30 minutes.)	weeks onthly aged 10 years and older with HeFH: weeks onthly ic patients aged 10 years and older with Ho onthly weeks ree 140mg/ml injections consecutively	FH:	
OTHER	□ 140mg/ml single-dose Prefilled SureClick® Autoinjector □ 420mg/3.5ml single-dose Pushtronex® system R SIGNATURE: I authorize pharmacy to act as my design	□ 140mg SQ every 2 □ 420mg SQ once m In pediatric patients: □ 140mg SQ every 2 □ 420mg SQ once m In adults and pediatri □ 420mg SQ once m □ 420mg SQ every 2 (For 420mg: Inject the within 30 minutes.) ee for initiating and coordination	weeks onthly aged 10 years and older with HeFH: weeks onthly ic patients aged 10 years and older with Ho onthly weeks ree 140mg/ml injections consecutively	FH:	